Application No. 09/782,588 Filed: February 12, 2001

Appendix A

**Pending Claims** 

1. A microscope slide composition comprising:

a) a substrate with a surface comprising discrete sites, said sites separated by a distance of less than 50 mm, wherein said substrate is formatted to the dimensions of

a microscope slide; and

b) a population of microspheres comprising at least a first and a second

subpopulation, wherein said first subpopulation comprises a first bioactve agent and

said second subpopulation comprises a second bioactive agent wherein said

microspheres are randomly distributed on said surface.

2. A composition according to claim 1, wherein said sites are separated by a distance

of less than 25 mm.

3. A composition according to claim 1, wherein said sites are separated by a distance

of less than 15 mm.

4. A composition according to claim 1, 2 or 3, wherein said sites are separated by a

distance of at least about 5 mm.

6. (Amended) The composition according to claim 1, wherein the distance between

centers of a first and second microsphere of said first subpopulation is at least 5 mm.

7. The composition according to claim 6, wherein the distance between said first and

-4-

Application No. 09/782,588 Filed: February 12, 2001

second microsphere of said first subpopulation is less than about 100 mm.

8. (Amended) A composition according to claim 1, wherein said substrate further comprises first and second assay locations, wherein said first and second subpopulations are distributed in said first and second assay locations.

- 9. A composition according to claim 8, wherein the distance between a first and second microsphere of said first subpopulation is less than about 100 mm.
- 10. A composition according to claim 9, wherein the distance between a first and second member of said first subpopulation is less than about 50 mm.
- 11. A composition according to claim 9, wherein the distance between a first and second member of said first subpopulation is less than about 15 mm.
- 12. A composition according to claim 9, 10 or 11, wherein the distance between said first and second member of said first subpopulation is at least about 5 mm.
- 18. A method for making a microscope slide composition comprising:
- a) providing a substrate with a surface comprising discrete sites, said sites separated by a distance of less than 50 mm, wherein said substrate is formatted to the dimensions of a microscope slide; and
- b) randomly distributing population of microspheres comprising at least a first and a second subpopulation, wherein said first subpopulation comprises a first

Application No. 09/782,588

Filed: February 12, 2001

bioactve agent and said second subpopulation comprises a second bioactive agent.

19. The method according to claim 18 wherein said wells are separated by a distance

of less than 25 mm.

20. The method according to claim 18, wherein said wells are separated by a distance

of less than 15 mm.

21. The method according to claim 18, wherein the ratio of said first and said second

subpopulation is at least 1: 36.

22. The method according to claim 18, wherein the ratio of said first and said second

subpopulation is at least 1: 100.

23. The method according to claim 18, wherein the distance between the centers a first

and second microsphere of said first subpopulation is at least 5 mm.

24. The method according to claim 18, wherein the distance between the centers of a

first and second microsphere of said first subpopulation is at least 15 mm.

25. The method according to claim 18, wherein the distance between a first and

second microsphere of said first subpopulation is at least 50 mm.

-6-

Application No. 09/782,588 Filed: February 12, 2001

Appendix B

Marked-up Version of the Claims

- 6. The composition according to claim 1 [or 5], wherein the distance between centers of a first and second microsphere of said first subpopulation is at least 5 mm.
- 8. A composition according to claim 1 [or 5], wherein said substrate further comprises first and second assay locations, wherein said first and second subpopulations are distributed in said first and second assay locations.